

Bayer CropScience LP P.O. Box 12014 2 T.W. Alexander Drive Research Triangle Park, North Carolina 27709 1-866-99BAYER (1-866-992-2937) ADMIRE[®] PRO Systemic Protectant For Post-Harvest or Dormant Treatment Use on Cranberry to control Black vine weevil.

EPA Reg. No. 264-827 SLN. No. OR-050025

24(c) Special Local Need

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF OREGON

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This label must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

This label for ADMIRE® PRO Systemic Protectant expires and must not be distributed or used in accordance with this SLN registration after December 31, 2020.

ADMIRE[®] PRO Systemic Protectant

ACTIVE INGREDIENT:	
Imidacloprid, 1-[(6-Chloro-3-pyridinyl)methyl]-N-nitro-2-imidazolidinimine	. 42.8%
OTHER INGREDIENTS:	. <u>57.2%</u>
Contains 4.6 pounds of active ingredient per gallon or 550 grams Al/liter	100.0%

SHAKE WELL BEFORE USING

KEEP OUT OF REACH OF CHILDREN CAUTION

For <u>MEDICAL</u> And <u>TRANSPORTATION</u> Emergencies <u>ONLY</u> Call 24 Hours A Day 1-800-334-7577 For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

ENVIRONMENTAL HAZARDS

Do not apply directly to water, areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. This product is toxic to wildlife and highly toxic to aquatic invertebrates.

This chemical demonstrates the properties and characteristics associated with chemicals detected in ground water. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

CRANBERRY

Pest Controlled	Rate fluid ounces/Acre
Black vine weevil	7.0

Post-Harvest or Dormant Spray

Apply specified rate of ADMIRE PRO to moist soil in one of the following methods:

- 1. As a broadcast soil spray (ground application only) or spot-spray directed to the infested bed area using a minimum of 20 gallons of water per acre followed **immediately** by 0.1 to 0.3 inches of rain or irrigation water. If making spot applications, use a rate per acre equivalent to a broadcast application.
- 2. As a chemigation application with 600 to 1000 gallons of water followed by 0.1 to 0.3 inches of irrigation water.

For best control of black vine weevil, application timing should target the larval stages and should be made only when crop is dormant (November 1 to March 1).

RESTRICTIONS

- Do not apply during bloom.
- Do not apply within 30 days of harvest.
- Do not make more than one post-harvest application in the same crop year.
- Do not make more than one dormant application in the same crop year.
- The main container label (Section 3) provides directions for post-bloom use to control rootgrubs and rootworms. This supplemental label provides directions for post-harvest and/or dormant use to control black vine weevil. Regardless of application timing, do not apply more than a total of 14 fluid ounces of ADMIRE PRO (0.5 lb. Imidacloprid) per acre per crop year.

NOTE: ADMIRE PRO has not been tested for crop response in tank mixes with other registered cranberry fungicides or insecticides. If tank mixing is desired, premix a sample of the ADMIRE PRO and the desired fungicide or insecticide partner at labeled rates and apply to a small area. Evaluate crop response within 48 hours and for at least two weeks prior to utilizing the tank mix on larger acreage. If crop injury results from the premix test, do not apply the tank mix to larger acreage.

NOTE: Follow these and all other Use Directions and Precautions on the package label.

For <u>MEDICAL</u> And <u>TRANSPORTATION</u> Emergencies <u>ONLY</u> Call 24 Hours A Day 1-800-334-7577 For <u>PRODUCT</u> <u>USE</u> Information Call 1-866-99BAYER (1-866-992-2937)

As with any crop-protection product, always read and follow label directions. For additional information call toll-free 1-866-99BAYER (1-866-992-2937).